arXiv:1603.03553v1 [math.AG] 11 Mar 2016

On a Projective Bundle Formula

James Fullwood[♠], Martin Helmer[♢]

[♠]Institute of Mathematical Research, The University of Hong Kong, Pok Fu Lam Road, Hong Kong.
[♢]Department of Mathematics, The University of California, Berkeley, USA.

Abstract

For a vector bundle $\mathscr{E} \to X$ with X a smooth variety over a field we prove a formula for the proper pushforward $\pi_* : A^*\mathbb{P}(\mathscr{E}) \to A^*X$, where $\pi : \mathbb{P}(\mathscr{E}) \to X$ denotes the associated projective bundle of *lines* in $\mathscr{E}$. Our formula is relative in the sense that it depends on the rank of $\mathscr{E}$ while being independent of the dimension of X. As applications, we apply the formula to compute the pushforward to X of the Chern class of a fibration $Y \to X$, whose fibers are degree d projective hypersurfaces with d arbitrary, and then derive a 'relative Bézout-type formula' for the intersection of $\dim(\mathbb{P}(\mathscr{E}))$ divisors in $\mathbb{P}(\mathscr{E})$ in terms of invariants of X.

Contents

1 Introduction **2**

2 Proof of Main Theorem **5**

3 Chern Class of a Fibration **8**

4 A Bézout Formula **13**

1 Introduction

Let X be a smooth variety over field $\mathfrak{K}$ endowed with a vector bundle $\mathscr{E} \to X$, and denote by $\pi : \mathbb{P}(\mathscr{E}) \to X$ the associated projective bundle of *lines* in $\mathscr{E}$. A fundamental task in the investigation of $\mathbb{P}(\mathscr{E})$ is to relate its invariants to invariants of X, the prototypical example of which being the formula for the Chow ring $A^*\mathbb{P}(\mathscr{E})$ in terms of A^*X, namely

$$A^*\mathbb{P}(\mathscr{E}) \cong A^*X[\zeta]/\left(\zeta^r + c_1(\mathscr{E})\zeta^{r-1} + \cdots + c_r(\mathscr{E})\right), \tag{1.1}$$

where $\zeta = c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1))$, $r = \mathrm{rk}(\mathscr{E})$ and $c_i(\mathscr{E})$ denotes the ith Chern class of $\mathscr{E}$. Other than its simplicity an attractive feature of formula (1.1) is that the RHS only depends on the rank of $\mathscr{E}$, thus being independent of the dimension of X.

In this note we consider the map

$$\pi_* : A^*\mathbb{P}(\mathscr{E}) \to A^*X,$$

i.e., the proper pushforward of algebraic cycles modulo rational equivalence associated with the bundle projection π, and derive a formula for π_* which (in a similar vein) only depends on the rank of $\mathscr{E}$. Our motivation for such a formula comes primarily from the fact that

$$\int_{\mathbb{P}(\mathscr{E})} \alpha = \int_X \pi_*\alpha \quad \forall \alpha \in A^*\mathbb{P}(\mathscr{E}), \tag{1.2}$$

so that invariants of $\mathbb{P}(\mathscr{E})$ (and of its closed subschemes) captured by the Chow ring may be expressed in terms of invariants of X, which are naturally much simpler. Note that the integral sign in (1.2) denotes the proper pushforward associated with the corresponding structure morphism to $\mathrm{spec}(\mathfrak{K})$. To state our formula we first introduce some notation.

Denote by $L_1, \ldots, L_m$ the *distinct* non-trivial Chern roots of $\mathscr{E}$ and let k_i be such that the multiplicity of L_i is $k_i + 1$, so that

$$c(\mathscr{E}) = (1 + L_1)^{k_1+1} \cdots (1 + L_m)^{k_m+1} \in A^*X.$$

We then consider the ring $A^*X[x_1, \ldots, x_m]$ where the x_i are formal variables, and let D be the operator on this ring given by

$$g \mapsto \frac{1}{k_1! \cdots k_m!} \frac{\partial^{k_1+\cdots+k_m}}{\partial x_1^{k_1} \cdots \partial x_m^{k_m}} (x_1^{k_1} \cdots x_m^{k_m} \cdot g).$$

Now let $\alpha \in A^*\mathbb{P}(\mathscr{E})$ be arbitrary, which by formula (1.1) may be written as

$$\alpha = \alpha_0 + \alpha_1\zeta + \alpha_2\zeta^2 + \cdots,$$

where $\alpha_j\zeta^j$ denotes $\pi^*\alpha_j \cdot \left(c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]\right)^j$ with $\alpha_j \in A^*X$. We then associate with α the elements

$$g_\alpha(x_i) = \left.\frac{\alpha - (\alpha_0 + \cdots + \alpha_{n-1}\zeta^{n-1})}{\zeta^{n-m+1}}\right|_{\zeta=x_i} \in A^*X[x_1, \ldots, x_m],$$

where $n = \mathrm{rk}(\mathscr{E}) - 1$. Our formula is then given by the following expression, which will be formally stated and proved as Theorem 2.1 below.

Theorem (Main Theorem)**.** *With notations as above, the map $\pi_* : A^*\mathbb{P}(\mathscr{E}) \to A^*X$ is given by*

$$\pi_*\alpha = \left.\left(D \cdot \sum_{i=1}^{m} \frac{g_\alpha(x_i)}{\prod_{l=1,\, l\neq i}^{m}(x_i - x_l)}\right)\right|_{x_1=-L_1,\ldots,x_m=-L_m} \in A_*X. \tag{1.3}$$

Clearly formula (1.3) is symmetric in the distinct Chern roots of $\mathscr{E}$ so that the end result may be expressed in terms of the Chern classes $c_i(\mathscr{E})$ and the α_j for $j \geq n$. The (non-invertible) terms $\prod_{l=1,\, l\neq i}^{m}(x_i - x_l)$ appearing in the denominator of the formula are formal expressions that end up cancelling before evaluation at the L_i. When $\mathscr{E}$ is the direct sum of tensor powers of a fixed line bundle on X we recover the main formula derived in [5].

For projective bundles of small rank, formula (1.3) is reasonable to apply by hand, though in any case a computer implementation is straightforward and is of negligible cost. We note that such an implementation may be of utility for software packages computing the characteristics classes of varieties. In particular an implementation of (1.3) allows the applicability setting of algorithms for computing singular Chern classes and Euler characteristics of (possibly singular) projective varieties and schemes (such as [9, 8]) to be broadened to the setting of projective bundles. To this end, and to simplify computations for the interested reader, we provide a test implementation in the form of a Macaulay2 [7] package which can be downloaded from `https://github.com/Martin-Helmer/PBF`.

For an explicit example of an application of the Main Theorem (proved as Theorem 2.1 below), let X be a smooth Fano variety over $\mathbb{C}$,

$$\mathscr{E} = \mathscr{O}_X \oplus \mathscr{O}(-2K_X) \oplus \mathscr{O}(-3K_X),$$

and let $\iota : Y \hookrightarrow \mathbb{P}(\mathscr{E})$ be a smooth anti-canonical hypersurface in $\mathbb{P}(\mathscr{E})$ given by the Weierstraß equation

$$Y : \left(y^2 = x^3 + fxz^2 + gz^3\right) \subset \mathbb{P}(\mathscr{E}),$$

where the coefficients f and g are sections of $\mathscr{O}(-4K_X)$ and $\mathscr{O}(-6K_X)$ respectively. The map $\varphi = \pi \circ \iota$ then endows Y with the structure of an elliptic fibration $\varphi : Y \to X$. Motivated by physical considerations, in [11] Sethi, Vafa and Witten derived a formula

for the Euler characteristic of Y in terms of the Chern classes of X for the case $\dim(X) = 3$, namely

$$\chi(Y) = \int_X 12c_1(X)c_2(X) + 360c_1(X)^3. \tag{1.4}$$

We note that the above formula may be simplified further since we necessarily have $c_1(X)c_2(X) = 24$ for X Fano. It immediately follows via Theorem 2.1 that formula (1.4) is simply a manifestation of formula (1.2) for $\alpha = \iota_* c(Y)$, i.e.,

$$\int_{\mathbb{P}(\mathscr{E})} \iota_* c(Y) = \int_X \pi_* \left(\iota_* c(Y)\right), \tag{1.5}$$

as the LHS of equation (1.5) coincides with $\chi(Y)$ by the Gauß-Bonnet theorem, and Theorem 2.1 immediately yields the RHS to be $\int_X 12c_1(X)c_2(X) + 360c_1(X)^3$.

More precisely, by the adjunction formula and standard exact sequences of vector bundles we have

$$\iota_* c(Y) = \frac{(1+\zeta)(1+\zeta-2K_X)(1+\zeta-3K_X)(3\zeta-6K_X)}{(1+3\zeta-6K_X)} \cdot \pi^* c(X),$$

where again $\zeta = c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]$. Thus

$$\begin{aligned} \varphi_* c(Y) &= \pi_*(\iota_* c(Y)) \\ &= \pi_* \left(\frac{(1+\zeta)(1+\zeta-2K_X)(1+\zeta-3K_X)(3\zeta-6K_X)}{(1+3\zeta-6K_X)} \right) c(X), \end{aligned}$$

where the second equality follows from the projection formula. We then take

$$\alpha = \frac{(1+\zeta)(1+\zeta-2K_X)(1+\zeta-3K_X)(3\zeta-6K_X)}{(1+3\zeta-6K_X)} = \alpha_0 + \alpha_1\zeta + \cdots \in A^*\mathbb{P}(\mathscr{E}),$$

where the α_i are the Taylor coefficients of α when expanded with respect to ζ. By Theorem 2.1 we then have

$$\begin{aligned} \pi_*\alpha &= \left(\frac{\alpha - (\alpha_0 + \alpha_1\zeta)}{\zeta(-3K_X + 2K_X)}\bigg|_{\zeta = 2K_X} \right) + \left(\frac{\alpha - (\alpha_0 + \alpha_1\zeta)}{\zeta(-2K_X + 3K_X)}\bigg|_{\zeta = 3K_X} \right) \\ &= \frac{-12K_X}{1-6K_X}, \end{aligned}$$

thus

$$\varphi_* c(Y) = \frac{12c_1(X)}{1+6c_1(X)} c(X). \tag{1.6}$$

Such a formula is often referred to in the literature as a *Sethi-Vafa-Witten formula* (see [5][1][4]). We note that one of the advantages of our methods is the dimension-independent

nature of Theorem 2.1, so that equation (1.6) is valid for X of any dimension. One may then obtain an analogue of formula (1.4) for $\chi(Y)$ with $\dim(X)$ arbitrary simply by reading off the term of codimension $\dim(X)$ on the RHS of equation (1.6), namely

$$\chi(Y) = \int_X 12c_1(X) \sum_{i=0}^{\dim(X)-1} c_i(X)\,(-6c_1(X))^{\dim(X)-1-i}\,.$$

In an analogous fashion one may use Theorem 2.1 to realize other invariants of closed subschemes of a projective bundle $\mathbb{P}(\mathscr{E})$ –such as intersection numbers, Chern numbers, Hirzebruch genera, etc.– in terms of invariants of its base, which as noted earlier are often much simpler. We devote §2 to the statement and proof of Theorem 2.1 which is the Main Theorem given in the introduction above; in the remaining sections we derive some applications of this theorem. In particular, we use Theorem 2.1 to derive a Bézout-type formula for the intersection of $\dim(\mathbb{P}(\mathscr{E}))$ divisors in $\mathbb{P}(\mathscr{E})$ in terms of the Chern classes of $\mathscr{E}$ and TX, and we generalize formula (1.6) to the case where Y is a general hypersurface in an arbitrary projective bundle $\mathbb{P}(\mathscr{E}) \to X$, so that Y is a general fibration of projective hypersurfaces.

Notation and conventions. The assumptions made at the outset of this section will hold throughout the rest of the paper. In particular, we will work over an arbitrary field $\mathfrak{K}$, except in §3 where we assume that $\mathfrak{K}$ is an algebraically closed field of characteristic zero. We denote by X an arbitrary smooth $\mathfrak{K}$-variety, which will always come endowed with a vector bundle $\mathscr{E} \to X$. We denote by $\pi : \mathbb{P}(\mathscr{E}) \to X$ the associated projective bundle of *lines* in $\mathscr{E}$. The *distinct* non-trivial Chern roots of $\mathscr{E}$ will be denoted by $L_1, ..., L_m$, and $k_i \in \mathbb{Z}$ will be such that the multiplicity of L_i is $k_i + 1$. We will always assume $m \leq \mathrm{rk}(\mathscr{E}) - 1$, as $\mathbb{P}(\mathscr{E}) \cong \mathbb{P}(\mathscr{E} \otimes \mathscr{L})$ for any line bundle $\mathscr{L}$ implies without loss of generality that we can set one of the Chern roots of $\mathscr{E}$ equal to zero.

2 Proof of Main Theorem

Let $\alpha \in A^*\mathbb{P}(\mathscr{E})$ be arbitrary, which by the isomorphism

$$A^*\mathbb{P}(\mathscr{E}) \cong A^*X[\zeta]/\left(\zeta^r + c_1(\mathscr{E})\zeta^{r-1} + \cdots + c_r(\mathscr{E})\right)$$

may be written as

$$\alpha = \alpha(\zeta) = \alpha_0 + \alpha_1\zeta + \alpha_2\zeta^2 + \cdots\,,$$

where we recall $\alpha_j\zeta^j$ denotes $\pi^*\alpha_j \cdot \left(c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]\right)^j$. We now prove the following

Theorem 2.1 (Formal Statement of Main Theorem)**.** *With notations as above we have that the map* $\pi_* : A^*\mathbb{P}(\mathscr{E}) \to A^*X$ *is given by*

$$\pi_*\alpha = \left(D \cdot \sum_{i=1}^{m} \frac{g_\alpha(x_i)}{\prod_{l=1,\, l\neq i}^{m}(x_i - x_l)} \right)\Bigg|_{x_1=-L_1,\ldots,x_m=-L_m} \in A^*X,$$

where

$$g_\alpha(x_i) = \frac{\alpha(x_i) - \left(\alpha_0 + \alpha_1 x_i + \cdots + \alpha_{\mathrm{rank}(\mathscr{E})-2} x_i^{\mathrm{rank}(\mathscr{E})-2}\right)}{x_i^{\mathrm{rank}(\mathscr{E})-m} \prod_{l=1,\, l\neq i}^{m}(x_i - x_l)},$$

and D *is the differential operator on* $A^*X[x_1, \ldots, x_m]$ *given by*

$$g \mapsto \frac{1}{k_1! \cdots k_m!} \frac{\partial^{k_1+\cdots+k_m}}{\partial x_1^{k_1} \cdots \partial x_m^{k_m}} (x_1^{k_1} \cdots x_m^{k_m} \cdot g).$$

Proof. As $\alpha \in A^*\mathbb{P}(\mathscr{E})$ may be written as

$$\alpha = \sum_{j=0} \pi^*\alpha_j \cdot \zeta^j,$$

by the projection formula for intersection products we have

$$\pi_*\alpha = \sum_{j=0} \alpha_j \cdot \pi_*(\zeta^j),$$

thus $\pi_*\alpha$ is completely determined by the $\pi_*(\zeta^j)$. For this, we use the fact that by Fulton's definition of Chern class ([6], §3.2) we have

$$\pi_*\left(1 + \zeta + \zeta^2 + \cdots\right) = c(\mathscr{E})^{-1} = \frac{1}{(1+L_1)^{k_1+1} \cdots (1+L_m)^{k_m+1}} \in A^*X, \tag{2.1}$$

where we recall that the L_i are the *distinct* non-trivial Chern roots of $\mathscr{E}$ and the k_i are uniquely determined non-negative integers. As the proper pushforward of a pure dimensional cycle (not in the kernel of π_*) preserves its dimension, it follows that $\pi_*(\zeta^j)$ coincides with the term of dimension $\dim(\mathbb{P}(\mathscr{E})) - j$ in the formal series expansion of $\left((1+L_1)^{k_1+1} \cdots (1+L_m)^{k_m+1}\right)^{-1}$ (so that $\pi_*(\zeta^j) = 0$ for $j < \mathrm{rk}(\mathscr{E}) - 1$). Now since

$$\frac{1}{(1+L_i)^{k_i+1}} = \frac{1}{k_i!} \frac{d^{k_i}}{dx_i^{k_i}} \left(\frac{1}{1-x_i}\right)\Bigg|_{x_i=-L_i} = \left(\sum_{j=0} \frac{(k_i+j)!}{k_i! j!} x^j\right)\Bigg|_{x_i=-L_i},$$

the term of codimension d in X of $c(\mathscr{E})^{-1}$ is precisely

$$\sum_{i_1+\cdots+i_m=d} \beta_{i_1}^{(k_1)}(-L_1)^{i_1}\cdot\beta_{i_2}^{(k_2)}(-L_2)^{i_2}\cdots\beta_{i_m}^{(k_m)}(-L_m)^{i_m},$$

where

$$\beta_{i_l}^{(k_l)} = \begin{cases} \frac{(k_l+i_l)!}{k_l!i_l!} & \text{for } i_l \neq 0 \\ 1 & \text{for } i_l = 0. \end{cases}$$

Now let $n = \mathrm{rk}(\mathscr{E})-1$. It then follows that $\dim(\zeta^n) = \dim(X)$, so that the term $\pi_*(\zeta^{n+j})$ is of codimension j in X. We then have

$$\pi_*(\zeta^{n+j}) = \begin{cases} 0 & \text{for} \quad j<0 \\ \displaystyle\sum_{i_1+\cdots+i_m=j} \beta_{i_1}^{(k_1)}(-L_1)^{i_1}\cdot\beta_{i_2}^{(k_2)}(-L_2)^{i_2}\cdots\beta_{i_m}^{(k_m)}(-L_m)^{i_m} & \text{for} \quad j\geq 0. \end{cases} \tag{2.2}$$

Thus

$$\begin{aligned}
\pi_*\alpha &= \pi_*\left(\sum_{j=-n}\alpha_{n+j}\zeta^{n+j}\right) \\
&= \sum_{j=-n}\alpha_{n+j}\cdot\pi_*\left(\zeta^{n+j}\right) \quad \text{(By the projection formula)} \\
&\overset{(2.2)}{=} \sum_{j=0}\alpha_{n+j}\left(\sum_{i_1+\cdots+i_m=j}\beta_{i_1}^{(k_1)}(-L_1)^{i_1}\cdot\beta_{i_2}^{(k_2)}(-L_2)^{i_2}\cdots\beta_{i_m}^{(k_m)}(-L_m)^{i_m}\right) \\
&= \sum_{j=0}\alpha_{n+j}\left(\sum_{i_1+\cdots+i_m=j}\frac{1}{k_1!}\frac{d^{k_1}}{dx_1^{k_1}}\left(x_1^{i_1+k_1}\right)\Big|_{x_1=-L_1}\cdots\frac{1}{k_m!}\frac{d^{k_m}}{dx_m^{k_m}}\left(x_m^{i_m+k_m}\right)\Big|_{x_m=-L_m}\right) \\
&= \sum_{j=0}\alpha_{n+j}\left(\frac{1}{k_1!\cdots k_m!}\frac{\partial^{k_1+\cdots+k_m}}{\partial x_1^{k_1}\cdots\partial x_m^{k_m}}\sum_{i_1+\cdots+i_m=j}x_1^{i_1+k_1}\cdots x_m^{i_m+k_m}\right)\Big|_{x_1=-L_1,\ldots,x_m=-L_m} \\
&= \sum_{j=0}\alpha_{n+j}\left(\frac{1}{k_1!\cdots k_m!}\frac{\partial^{k_1+\cdots+k_m}}{\partial x_1^{k_1}\cdots\partial x_m^{k_m}}x_1^{k_1}\cdots x_m^{k_m}\sum_{i_1+\cdots+i_m=j}x_1^{i_1}\cdots x_m^{i_m}\right)\Big|_{x_1=-L_1,\ldots,x_m=-L_m} \\
&\overset{(2.3)}{=} \sum_{j=0}\alpha_{n+j}\left(D\cdot\sum_{i=1}^m\frac{x_i^{m+j-1}}{\prod_{l=1,\,l\neq i}^m(x_i-x_l)}\right)\Big|_{x_1=-L_1,\ldots,x_m=-L_m} \\
&= \left(D\cdot\sum_{i=1}^m\frac{\sum_{j=0}\alpha_{n+j}x_i^{m+j-1}}{\prod_{l=1,\,l\neq i}^m(x_i-x_l)}\right)\Big|_{x_1=-L_1,\ldots,x_m=-L_m} \\
&= \left(D\cdot\sum_{i=1}^m\frac{\alpha(x_i)-\left(\alpha_0+\alpha_1x_i+\cdots+\alpha_{\mathrm{rank}(\mathscr{E})-2}x_i^{\mathrm{rank}(\mathscr{E})-2}\right)}{x_i^{\mathrm{rank}(\mathscr{E})-m}\prod_{l=1,\,l\neq i}^m(x_i-x_l)}\right)\Bigg|_{\substack{x_j=-L_j,\\ j=1,\ldots,m}}.
\end{aligned}$$

Note that in the seventh equality we used the fact that (see [3])

$$\sum_{i_1+\cdots+i_m=j} x_1^{i_1}x_2^{i_2}\cdots x_m^{i_m} = \sum_{i=1}^{m} \frac{x_i^{m+j-1}}{\prod_{l=1,\, l\neq i}^{m}(x_i - x_l)}. \tag{2.3}$$

This concludes the proof of the theorem. □

Remark 2.1. It follows from formula (2.3) (along with degree considerations) that terms of the form $L_i^{\mathrm{rank}(\mathscr{E})-m}\prod_{l=1,\, l\neq i}^{m}(L_i - L_l)$ appearing in final expression for $\pi_*\alpha$ in fact cancel once put over a common denominator. As such, in practice one may in fact avoid the use of the differential operator D by first assuming *all* the non-trivial Chern roots are distinct, and then evaluating the final simplified rational expression at the possibly non-distinct total number of non-trivial Chern roots of $\mathscr{E}$.

3 Chern Class of a Fibration

We first review the theory of Chern classes from a functorial perspective (for more details one may consult [2][12]). In this section we impose the constraint that $\mathfrak{K}$ denotes an algebraically closed field of characteristic zero. For V a $\mathfrak{K}$-variety denote by $K_0(\mathrm{Var}_V)$ the free $\mathbb{Z}$-module generated by isomorphism classes of proper morphisms to V, modulo the relation

$$[Y \xrightarrow{\varphi} V] = [Z \xrightarrow{\varphi|_Z} V] + [U \xrightarrow{\varphi|_U} V],$$

whenever Z is a closed subvariety of Y with open complement U in Y. We may endow $K_0(\mathrm{Var}_V)$ with a ring structure by setting

$$[Y \xrightarrow{\varphi} V]\cdot[W \xrightarrow{\psi} V] = [Y \times_V W \to V],$$

where $Y \times_V W \to V$ denotes the fibered product of Y with W over V (with respect to the morphisms φ and ψ). We refer to this ring $K_0(\mathrm{Var}_V)$ as the *relative Grothendieck ring of varieties* (over V). When $V = \mathrm{spec}(\mathfrak{K})$ the relative Grothendieck ring is often denoted $K_0(\mathrm{Var}_{\mathfrak{K}})$, and referred to simply as the *Grothendieck ring of varieties*. Moreover, we may view $K_0(\mathrm{Var}_{(\cdot)})$ as a covariant functor from varieties to groups with respect to proper morphisms.

For this, given a proper morphism $V \xrightarrow{f} S$ we define a homomorphism

$$K_0(f) : K_0(\mathrm{Var}_V) \longrightarrow K_0(\mathrm{Var}_S)$$

given by

$$[Y \to V] \mapsto [Y \to V \to S].$$

With these prescriptions it is straightforward to show that indeed $K_0(\mathrm{Var}_{(\cdot)})$ is a covariant functor with respect to proper morphisms, which we refer to as the *relative Grothendieck functor*.

Now denote by A_* the *Chow group functor*, which takes a variety V to its group of algebraic cycles modulo rational equivalence A_*V, and acts on proper morphisms via proper pushforward of algebraic cycles. For V smooth the Chow group may be given a ring structure via Fulton's intersection products [6]. In such a case we will denote A_*V by A^*V, which we refer to as the Chow *ring* of V (see [6], § 1.4.). For a proper morphism $f : V \to S$ the associated pushforward is denoted $f_* : A_*V \to A_*S$. The Chow group functor is also covariant with respect to proper morphisms, and there exists a unique natural transformation

$$c_* : K_0(\mathrm{Var}_{(\cdot)}) \longrightarrow A_*$$

such that if V is smooth then

$$c_*(V)([V \xrightarrow{\mathrm{id}} V]) = c(TV) \cap [V] \in A^*V,$$

where $c_*(V)$ denotes the homomorphism $c_*(V) : K_0(\mathrm{Var}_V) \to A_*V$ induced by the natural transformation c_*, and $c(TV) \cap [V]$ denotes the total homological Chern class of V (again see [2][12] for details). For V possibly singular the class $c_*(V)([V \xrightarrow{\mathrm{id}} V])$ will be denoted by $c_{\mathrm{SM}}(V)$, which is referred to in the literature as the *Chern-Schwartz-MacPherson class* of V. It follows from functoriality that for a more general morphism $\varphi : Y \to V$ we have

$$c_*(V)([Y \xrightarrow{\varphi} V]) = \varphi_* c_{\mathrm{SM}}(Y),$$

which from here on will be denoted by $c_{\mathrm{SM}}(\varphi)$, and referred to as the Chern-Schwartz-MacPherson class of the *morphism* φ. If Y and V are both smooth then we will simply refer to $c_{\mathrm{SM}}(\varphi)$ as the Chern class of φ, written $c(\varphi)$, so that the usual Chern class of a smooth variety may be referred to as the Chern class of the identity morphism.

From this perspective we may reformulate the Sethi-Vafa-Witten formula (1.6) from §1 as

$$c(\varphi) = \frac{12c_1(X)}{1 + 6c_1(X)} \cdot c(\mathrm{id}_X) \in A^*X. \tag{3.1}$$

We now generalize this formula to the case where Y is an arbitrary smooth hypersurface in a projective bundle, not necessarily an elliptic fibration.

So let X be a smooth $\mathfrak{K}$-variety endowed with a vector bundle $\mathscr{E} \to X$ (as in the Notation and Conventions stated in §1). Assume Y is the smooth zero-scheme of a non-trivial section of a line bundle $\mathscr{M} \to \mathbb{P}(\mathscr{E})$, so that Y is an embedded smooth hypersurface in $\mathbb{P}(\mathscr{E})$. Denote by $\iota : Y \hookrightarrow \mathbb{P}(\mathscr{E})$ the natural inclusion, so that $\varphi = \pi \circ \iota : Y \to X$ endows Y with the structure of a fibration of degree d hypersurfaces in $\mathbb{P}^{\mathrm{rk}(\mathscr{E})-1}$, where d is such that

$[Y] = d\zeta + \pi^*\beta$ with β a divisor in X, and $\zeta = c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]$ ($[Y]$ is of this form by Theorem 3.3 of [6]). We then associate with Y a rational expression Q_Y in the distinct non-trivial Chern roots L_i of $\mathscr{E}$ and the class $\beta \in A^1(X)$. Note that the expression Q_Y will be seen to arise from the application of Theorem 2.1 to the current setting in proof of Theorem 3.1 below.

To define Q_Y, we first consider the the following sum in the formal variables x_j and β

$$\sum_{i=1}^{n} \left(\frac{(1-x_i)\prod_{j=1}^{n}(1+x_j-x_i)}{x_i \prod_{j=1,\, j\neq i}^{n}(x_i - x_j)} \cdot \frac{\beta - d\cdot x_i}{1+\beta - d\cdot x_i} - \frac{\prod_{j=1}^{n}(1+x_j)}{x_i \prod_{j=1,\, j\neq i}^{n}(x_i - x_j)} \cdot \frac{\beta}{1+\beta} \right),$$

where $n = \mathrm{rk}(\mathscr{E}) - 1$, which coincides with the *total* number of non-trivial Chern roots of $\mathscr{E}$ by assumption. Then after putting the above sum over a common denominator, cancelling common factors from the numerator and denominator of the form $x_i \prod_{j=1,\, j\neq i}^{r}(x_i - x_j)$ (which one may in fact do as mentioned in Remark 2.1), and then evaluating the x_i at the n number of non-trivial Chern roots of $\mathscr{E}$ we arrive at an expression of the form

$$\frac{P_Y}{(1+\beta)\cdot(1+\beta - dL_1)^{k_1+1}\cdots(1+\beta-dL_m)^{k_m+1}},$$

where we recall $k_i + 1$ is the multiplicity of L_i, and P_Y is a polynomial in the non-trivial Chern roots of $\mathscr{E}$ and β. We then take

$$Q_Y = \frac{P_Y}{(1+\beta)\cdot(1+\beta - dL_1)^{k_1+1}\cdots(1+\beta-dL_m)^{k_m+1}}.$$

For example if Y is the elliptic fibration in Weierstrass form defined in §1 we have

$$Q_Y = \frac{12c_1(X)(1-3c_1(X))}{(1+6c_1(X))(1+6c_1(X)-6c_1(X))(1-3c_1(X))} = \frac{12c_1(X)}{1+6c_1(X)} \in A^*X.$$

We note that if $\mathscr{E}_d^\vee$ denotes a vector bundle whose distinct non-trivial Chern roots are $-dL_1, ..., -dL_m$, where $-dL_i$ has the same multiplicity as L_i (i.e., $k_i + 1$), then the denominator of Q_Y may be associated with the inverse Chern class of the vector bundle $\mathscr{E}_d^\vee \otimes \mathscr{O}(\beta)$, so that Q_Y may be written as

$$Q_Y = c(\mathscr{E}_d^\vee \otimes \mathscr{O}(\beta))^{-1} \cap P_Y \in A^*X.$$

The Chern class of φ is then given by the following

Theorem 3.1. *Let $\varphi : Y \to X$ be a fibration of degree d projective hypersurfaces as defined above. Then*

$$c(\varphi) = Q_Y \cdot c(\mathrm{id}_X) \in A^*X. \tag{3.2}$$

We first prove a lemma which will be useful not only for the proof of Theorem 3.1 but also in the next section for the proof of Theorem 4.1, namely

Lemma 3.2. *Let R be a commutative ring with unity and let $m \in \mathbb{N}$ be a positive integer. Then for any choice of $\alpha_0, \alpha_1, \ldots, \alpha_{m-1} \in R$ we have*

$$\sum_{i=1}^{m} \frac{\alpha_0 + \alpha_1 x_i + \cdots + \alpha_{m-1} x_i^{m-1}}{x_i \prod_{j=1,\, j\neq i}^{m} (x_i - x_j)} = \sum_{i=1}^{m} \frac{\alpha_0}{x_i \prod_{j=1,\, j\neq i}^{m} (x_i - x_j)} \in R(x_1, ..., x_m). \tag{3.3}$$

Proof. The LHS of equation (3.3) may be expanded as

$$\sum_{i=1}^{m} \alpha_0 \frac{x_i^{-1}}{\prod_{j=1,\, j\neq i}^{m} (x_i - x_j)} + \alpha_1 \sum_{i=1}^{m} \frac{1}{\prod_{j=1,\, j\neq i}^{m} (x_i - x_j)} + \cdots + \alpha_{m-1} \sum_{i=1}^{m} \frac{x_i^{m-2}}{\prod_{j=1,\, j\neq i}^{m} (x_i - x_j)},$$

and in (i) of the proof of Theorem 3.2 of [3] it is shown that for $0 \leq p \leq m-2$ we have

$$\sum_{i=1}^{m} \frac{x_i^p}{\prod_{j=1,\, j\neq i}^{m} (x_i - x_j)} = 0,$$

from which the lemma follows. □

Proof of Theorem 3.1. By Remark 2.1 it suffices to prove the case where the non-trivial Chern roots of $\mathscr{E}$ are all distinct, i.e., when $m = n = \mathrm{rk}(\mathscr{E}) - 1$. So let $\mathscr{O}(Y) \to \mathbb{P}(\mathscr{E})$ denote the line bundle corresponding to the divisor class of Y (whose restriction to Y yields the normal bundle to Y in $\mathbb{P}(\mathscr{E})$). In order to write an explicit expression for $c(\varphi)$, we first write an expression for $c(\iota)$, where we recall $\iota : Y \to \mathbb{P}(\mathscr{E})$ is the natural inclusion. For this, we consider the following exact sequences of vector bundles (as Chern classes are multiplicative)

$$0 \to TY \to \iota^* T\mathbb{P}(\mathscr{E}) \to \iota^* \mathscr{O}(Y) \to 0$$

$$0 \to T_{\mathbb{P}(\mathscr{E})/X} \to T\mathbb{P}(\mathscr{E}) \to \pi^* TX \to 0$$

$$0 \to \mathscr{O}_{\mathbb{P}(\mathscr{E})} \to \pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1) \to T_{\mathbb{P}(\mathscr{E})/X} \to 0.$$

The first sequence (from the top) is standard, the second one may take as the definition of the relative tangent bundle $T_{\mathbb{P}(\mathscr{E})/X}$, and the third is lifted from B.5.8 in [6]. We then

have

$$\begin{aligned}
c(\iota) &= \iota_* \left(c(TY) \cap [Y] \right) \\
&= \iota_* \left(\frac{c(\iota^* T\mathbb{P}(\mathscr{E}))}{c(\iota^* \mathscr{O}(Y))} \cap [Y] \right) \\
&= \frac{c(\pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cdot c(\pi^* TX)}{c(\mathscr{O}(Y))} \cap \iota_*[Y] \\
&= \frac{c(\pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cdot c(\pi^* TX)}{c(\mathscr{O}(Y))} \cap (c_1(\mathscr{O}(Y)) \cap \pi^*[X]) \\
&= \frac{c(\pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cdot c_1(\mathscr{O}(Y))}{c(\mathscr{O}(Y))} \cap (c(\pi^* TX) \cap \pi^*[X]) \\
&= \frac{c(\pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cdot c_1(\mathscr{O}(Y))}{c(\mathscr{O}(Y))} \cap \pi^* c(\mathrm{id}_X),
\end{aligned}$$

where the third equality follows from the projection formula, the fourth from the fact that $\pi^*[X] = [\mathbb{P}(\mathscr{E})]$ (since $\iota_*[Y] = c_1(\mathscr{O}(Y)) \cap [\mathbb{P}(\mathscr{E})]$), the fifth from the commutativity of Chern classes, and the sixth by the behavior of Chern classes with respect to pullback.

Now since $(1, L_1, ..., L_n)$ are the Chern roots of $\mathscr{E}$ we have that the Chern roots of $\pi^*(\mathscr{E}) \otimes \mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)$ are then $(\zeta, L_1 + \zeta, \ldots, L_n + \zeta)$, with $\zeta = c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]$ (note that technically, the Chern roots are of the form $\pi^* L_i + \zeta$ but we have omitted the pullbacks; we shall also slightly abuse notation and let $1 + Y$ denote $1 + c_1(Y)$). Thus

$$c(\iota) = \frac{(1+\zeta)(1+\zeta+L_1) \cdots (1+\zeta+L_n) \cdot [Y]}{1+Y} \cdot \pi^* c(\mathrm{id}_X),$$

so that

$$\begin{aligned}
c(\varphi) &= \varphi_* c(\mathrm{id}_Y) \\
&= \pi_* \circ \iota_* c(\mathrm{id}_Y) \\
&= \pi_* c(\iota) \\
&= \pi_* \left(\frac{(1+\zeta)(1+\zeta+L_1) \cdots (1+\zeta+L_n) \cdot [Y]}{1+Y} \cdot \pi^* c(\mathrm{id}_X) \right) \\
&= \pi_* \left(\frac{(1+\zeta)(1+\zeta+L_1) \cdots (1+\zeta+L_n) \cdot [Y]}{1+Y} \right) \cdot c(\mathrm{id}_X),
\end{aligned}$$

where the second equality follows from functoriality and the fifth follows via the projection formula. The theorem then follows once we show

$$\pi_* \left(\frac{(1+\zeta)(1+\zeta+L_1) \cdots (1+\zeta+L_n) \cdot [Y]}{1+Y} \right) = Q_Y.$$

So let

$$\alpha = \alpha(\zeta) = \frac{(1+\zeta)\cdot\prod_{i=1}^{n}(1+\zeta+L_i)\cdot[Y]}{1+Y}.$$

Applying Theorem 2.1 along with Lemma 3.2 to the class α then yields

$$\begin{aligned}\pi_*\alpha &= \sum_{i=1}^{m}\left(\frac{\alpha(x_i)-\alpha_0}{x_i\prod_{j=1,\,j\neq i}^{m}(x_i-x_j)}\right)\Bigg|_{x_1=-L_1,\ldots,x_m=-L_m} \\ &= \sum_{i=1}^{m}\left(\frac{(1-L_i)\prod_{j=1}^{m}(1+L_j-L_i)}{L_i\prod_{j=1,\,j\neq i}^{m}(L_j-L_i)}\cdot\frac{\beta-dL_i}{1+\beta-dL_i}-\frac{\prod_{j=1}^{m}(1+L_j)}{L_i\prod_{j=1,\,j\neq i}^{m}(L_j-L_i)}\cdot\frac{\beta}{1+\beta}\right),\end{aligned}$$

which coincides with Q_Y by definition, thus concluding the proof. □

In Table 3.1 we list Q_Y for various explicit examples.

Non-trivial Chern roots of $\mathscr{E}$	$[Y]$	Generic fiber	Q_Y
$(2L, 3L)$	$3\zeta + 6L$	Elliptic curve	$\frac{12L}{1+6L}$
(L, S)	$3\zeta + 2L + S$	Elliptic curve	$\frac{6(2L+2L^2-LS-S^2)}{(1+2L-2S)(1+2L+S)}$
$(L, 2L)$	$3\zeta + 4L$	Elliptic curve	$\frac{12L}{1+4L}$
(L, L, L)	$3\zeta + 3L$	del Pezzo surface	$\frac{3L+9}{1+3L}$
(L, L, L)	$4\zeta + 4L$	K3 surface	$\frac{24-12L}{1+4L}$
(L, L, L)	$5\zeta + 5L$	Surface of general-type	$\frac{55-45L}{1+5L}$
(L, L, L, L)	$5\zeta + 5L$	Calabi-Yau threefold	$\frac{280L-200}{1+5L}$

Table 3.1: Explicit examples of Q_Y

4 A Bézout Formula

An immediate corollary of Theorem 2.1 yields a formula for the intersection of $\dim(\mathbb{P}(\mathscr{E}))$ divisors in $\mathbb{P}(\mathscr{E})$ in terms of invariants of its base X, which we think of as a 'relative Bézout-type formula'. The formula is given in terms of elementary symmetric polynomials and complete homogeneous polynomials, which we now define.

We denote the j^{th} elementary symmetric polynomial in the p variables $y_1, \ldots, y_p$ by $e_j(y_1, \ldots, y_p)$, which is given by

$$e_j(y_1, \ldots, y_p) = \sum_{l_1 \leq l_2 \leq \cdots \leq l_j \leq p} y_{l_1} \cdots y_{l_j}. \tag{4.1}$$

The complete homogeneous symmetric polynomial of degree d in the q variables $y_1, \ldots, y_q$ will be denoted by $h_d(y_1, \ldots, y_q)$, and is given by

$$h_d(y_1, \ldots, y_q) = \sum_{\substack{l_1+\cdots+l_q=d \\ l_i \geq 0}} y_1^{l_1} \cdots y_q^{l_q}. \tag{4.2}$$

For more on the theory of such polynomials see for example [10] or [3].

We denote by $h_d(\mathscr{E})$ the dth homogeneous symmetric polynomial in the negative distinct non-trivial Chern roots of $\mathscr{E}$, i.e.

$$h_d(\mathscr{E}) = h_d(-L_1, \ldots, -L_m).$$

Before stating and proving our relative Bézout formula we recall that by Theorem 3.3 of [6] a divisor $D \in A^1\mathbb{P}(\mathscr{E})$ is of the form

$$D = \pi^* \underline{D} + d\zeta,$$

where $\underline{D}$ is a divisor in X, $\zeta = c_1(\mathscr{O}_{\mathbb{P}(\mathscr{E})}(1)) \cap [\mathbb{P}(\mathscr{E})]$ and d is an integer. The relative Bézout-type formula is then given by the following

Theorem 4.1. *Let $N = \dim(\mathbb{P}(\mathscr{E}))$, $D_1, \ldots, D_N$ be N distinct divisors in $\mathbb{P}(\mathscr{E})$, and let $\underline{D}_i$ and d_i be such that $D_i = \pi^*\underline{D}_i + d_i\zeta$. Denote by α the intersection product*

$$\alpha = D_1 \cdots D_N \in A^*\mathbb{P}(\mathscr{E}),$$

and denote by $e_j(\alpha)$ the jth elementary symmetric polynomial in $\underline{D}_i/d_i$, so that

$$e_j(\alpha) = e_j(\underline{D}_1/d_1, \ldots, \underline{D}_N/d_N).$$

Then

$$\pi_*\alpha = d_1 \cdots d_N \cdot \sum_{j=0}^{\dim(X)} h_{\dim(X)-j}(\mathscr{E}) e_j(\alpha) \in A^*X. \tag{4.3}$$

Proof. As in the proof of Theorem 3.1 we first assume all the non-trivial Chern roots of $\mathscr{E}$ are distinct, which by Remark 2.1 we may do without any loss of generality. Expanding α we obtain

$$\alpha = \alpha(\zeta) = d_1 \cdots d_N \sum_{j=0}^{N} \zeta^{N-j} e_j(\alpha).$$

Now let $n = \mathrm{rk}(\mathscr{E}) - 1$. Theorem 2.1 along with Lemma 3.2 then yields

$$\begin{aligned}\pi_*\alpha &= d_1 \cdots d_N \sum_{i=1}^{n} \left(\frac{\alpha(x_i) - d_1 \cdots d_N \cdot e_N(\alpha)}{x_i \prod_{j=1,\, j\neq i}^{n} (x_i - x_j)} \right) \Bigg|_{x_1 = -L_1, \ldots, x_n = -L_n} \\ &= d_1 \cdots d_N \sum_{i=1}^{n} \left(\frac{\sum_{j=0}^{N-1} x_i^{N-j} e_j(\alpha)}{\prod_{j=1,\, j\neq i}^{n} (x_i - x_j)} \right) \Bigg|_{x_1 = -L_1, \ldots, x_n = -L_n} .\end{aligned}$$

Expanding the summation, applying again Lemma 3.2 and recalling that $N = \dim(X) + n$ we obtain

$$\pi_*\alpha = d_1 \cdots d_N \cdot \sum_{j=0}^{dim(X)} h_{\dim(X)-j}(\mathscr{E}) e_j(\alpha),$$

as desired. □

References

[1] Paolo Aluffi and Mboyo Esole. Chern class identities from tadpole matching in type IIB and F-theory. *J. High Energy Phys.*, (3):032, 29, 2009.

[2] Jean-Paul Brasselet, Jörg Schürmann, and Shoji Yokura. Hirzebruch classes and motivic Chern classes for singular spaces. *J. Topol. Anal.*, 2(1):1–55, 2010.

[3] EF Cornelius Jr. Identities for complete homogeneous symmetric polynomials. *JP J. Algebra Number Theory Appl*, 21:109–116, 2011.

[4] Mboyo Esole, James Fullwood, and S.-T. Yau. D5 elliptic fibrations: Non-Kodaira fibers and new orientifold limits of F-theory. *arXiv:1110.6177*, 2011.

[5] James Fullwood. On generalized Sethi-Vafa-Witten formulas. *J. Math. Phys.*, 52(8):082304, 10, 2011.

[6] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge.* Springer-Verlag, Berlin, second edition, 1998.

[7] Daniel R. Grayson and Michael E. Stillman. *Macaulay2, a software system for research in algebraic geometry*, 2016.

[8] Martin Helmer. Algorithms to compute the topological Euler characteristic, Chern-Schwartz-Macpherson class and Segre class of projective varieties. *Journal of Symbolic Computation*, 2015.

[9] Christine Jost. An algorithm for computing the topological Euler characteristic of complex projective varieties. *arXiv:1301.4128*, 2013.

[10] Ian Grant Macdonald. *Symmetric functions and Hall polynomials*. Oxford university press, 1998.

[11] S. Sethi, C. Vafa, and E. Witten. Constraints on low-dimensional string compactifications. *Nuclear Phys. B*, 480(1-2):213–224, 1996.

[12] Shoji Yokura. Motivic characteristic classes. In *Topology of stratified spaces*, volume 58 of *Math. Sci. Res. Inst. Publ.*, pages 375–418. Cambridge Univ. Press, Cambridge, 2011.